

***Minimelanolocus atlanticus* sp. nov. and *M. navicularis* from the Brazilian Atlantic Forest**

PHELIPE M.O. COSTA¹, ELAINE MALOSSO^{1*}, MARCELA A. BARBOSA²,
WANDERSON L. TAVARES¹ & RAFAEL F. CASTAÑEDA-RUIZ³

¹ Centro de Biociências, Departamento de Micologia &

² Programa de Pós-Graduação em Biologia de Fungos,
Universidade Federal de Pernambuco, Avenida da Engenharia,
s/n Cidade Universitária, Recife, PE, 50.740-600, Brazil

³ Instituto de Investigaciones Fundamentales en Agricultura Tropical (INIFAT)
Alejandro de Humboldt, OSDE, Grupo Agrícola,
Calle 1 Esq. 2, Santiago de Las Vegas, C. Habana, Cuba, C.P. 17200

* CORRESPONDENCE TO: elaine.malosso@ufpe.br

ABSTRACT—*Minimelanolocus navicularis* and a new species, *M. atlanticus*, are described and illustrated from collections on decaying leaves of unidentified plants. *Minimelanolocus atlanticus* is distinguished by oblong, 2-septate conidia, with brown basal and central cells and gray to subhyaline apical cell and slightly flattened or sub-rounded apex. Notes are provided on *M. navicularis* recorded from Brazilian Atlantic Forest.

KEY WORDS—conidial fungi, asexual fungi, neotropic, systematic

Introduction

The genus *Minimelanolocus*, established by Castañeda-Ruiz et al. (2001) and typified by *Pseudospiropes navicularis* R.F. Castañeda [$\equiv$ *M. navicularis*], is characterized by conidiophores that are macronematous, mononematous, erect, unbranched, smooth or verrucose and conidiogenous cells that are polyblastic, sympodial elongated, terminal, indeterminate, and with inconspicuous or slightly prominent, narrow, opaque, refractive to somewhat obscure conidiogenous loci. *Minimelanolocus* currently comprises 31 species (Castañeda-Ruiz et al. 2001, 2003; Fiuza et al. 2017; Heredia et al. 2014; Hernández-Restrepo et al. 2012; Liu et al. 2015; Ma et al. 2008, 2011; Tian

et al. 2016; Xia et al. 2014; Zhang et al. 2009, 2010), most of which occur as saprobes on decaying leaves, rotten wood, dead branches, bamboo culms, and submerged plant debris in aquatic habitats.

During a mycological survey of microfungi associated with leaf litter in a Brazilian Atlantic forest, two conspicuous fungi representing *Minimelanolocus* were collected. We describe one as a new species and present a new description of the other, *M. navicularis*.

Materials & methods

Individual collections were placed in plastic bags, taken to the laboratory, and treated according to Castañeda-Ruiz et al. (2016). Mounts were prepared in lactic acid (90%) or in polyvinyl alcohol-glycerol (8 g PVA in 100 ml H₂O + 5 ml glycerol) and lactofuchsin (0.1 g acid fuchsin in 100 ml 85% lactic acid) following Carmichael (1955). Measurements were made at a magnification of $\times 1000$ under a Nikon Eclipse Ni-U microscope with bright field optics, and photomicrographs were taken using DIC optics with a Nikon DS-Fi2 camera. The holotype was deposited in the Herbarium of Universidade Federal de Pernambuco, Recife, Brazil (URM).

Taxonomy

Minimelanolocus atlanticus P.M.O. Costa, Malosso &

R.F. Castañeda, **sp. nov.**

FIGS 1, 2

INDEX FUNGORUM IF 554056

Differs from *Minimelanolocus obscurus* by its 2-septate and versicolored conidia.

TYPE: Brazil, Pernambuco, Igarassú, RPPN Refúgio Ecológico Charles Darwin, 7°48'S 34°57'W, on decaying leaves of an unidentified plant, 12.V.2017, coll. P.M.O. Costa (**Holotype**, URM 90133).

ETYMOLOGY: Latin, *atlanticus*, refers to habitat of the fungus, Brazilian Atlantic Forest.

COLONIES on the natural substrate, effuse, hairy, brown. Mycelium superficial and immersed, composed of septate, branched, brown, smooth-walled hyphae, 2–3 μm diam. CONIDIOPHORES macronematous, mononematous, erect, straight or flexuous, cylindrical, slightly geniculate near the apex, unbranched, 7–15-septate, dark brown below, pale brown toward the apex, 200–300 $\times$ 5–7 μm , smooth. CONIDIOGENOUS CELLS polyblastic, somewhat sinuate with inconspicuous denticulate or faintly cicatrized, opaque loci, terminal, integrated, indeterminate with several sympodial extensions, pale brown, 30–60 $\times$ 3.5–4 μm . Conidial secession schizolytic. CONIDIA solitary, acropleurogenous, oblong, sub-rounded or slightly flattened at the apex, 2-septate, bicolor with basal and central cells brown to dirty brown, apical cell smoked-gray to subhyaline, smooth, dry, 18–21 $\times$ 4–5 μm . Sexual state unknown.




FIG. 1. *Minimelanolocus atlanticus* (ex holotype URM 90133). A. Conidia; B. Conidiophores and conidiogenous cells; C. Conidiogenous cells; D. Conidiogenous cells and conidia. C & D share the same scale.




FIG. 2. *Minimelanolocus atlanticus* (ex holotype URM 90133).
Conidiogenous cells and conidia.

NOTES: *Minimelanolocus obscurus* (Matsush.) R.F. Castañeda & Heredia [= *Pseudospiropes obscurus* Matsush.] is similar to *M. atlanticus* in producing oblong conidia but is distinguished by conidia that are 1–4-septate (usually 3-septate), larger ($18\text{--}34 \times 4.5\text{--}8\text{ }\mu\text{m}$), and uniformly colored subhyaline or pale brown (Castañeda-Ruiz et al. 2001, Liu et al. 2015, Matsushima 1983).

Minimelanolocus navicularis (R.F. Castañeda) R.F. Castañeda,
Cryptog. Mycol. 22: 9 (2001).

FIG. 3

CONIDIOPHORES macronematous, mononematous, erect, straight or flexuous, unbranched, cylindrical, inflated at the base, 2–5-septate, smooth, brown, pale brown above, smooth, $43\text{--}70 \times 4\text{--}6\text{ }\mu\text{m}$. CONIDIOGENOUS CELLS polyblastic, integrated, terminal, indeterminate, sympodial elongated,




FIG. 3. *Minimelanolocus navicularis* (ex URM 90185).
A. Conidia; B. Conidiophores and conidiogenous cells.

inconspicuous denticulate, slightly melanized at the loci, $14\text{--}22 \times 3\text{--}4\text{ }\mu\text{m}$, pale brown. Conidial secession schizolytic. CONIDIA solitary, acropleurogenous, navicular, straight, smooth, 3-euseptate, $17\text{--}24 \times 5\text{--}8.5\text{ }\mu\text{m}$, obscure at the septa, dry, with subhyaline to pale brown end cells, brown to dark brown central cells.

SPECIMEN EXAMINED: **BRAZIL**, **PERNAMBUCO STATE**, Lagoa da Mata, Ilha de Itamaracá, $7^{\circ}46'S\ 34^{\circ}51'W$, on decaying leaves of an unidentified plant, 23.XI.2016, coll. W.L. Tavares (URM 90185).

NOTES: *Minimelanolocus navicularis* has been previously recorded in Brazil by Santa Izabel et al. (2011); their specimen had longer conidiophores ($45\text{--}155\text{ }\mu\text{m}$) than our specimen, but the conidia in both specimens closely match those of the type species ($20\text{--}25 \times 6\text{--}8\text{ }\mu\text{m}$; Castañeda-Ruiz 1987).

Acknowledgments

The authors express their sincere gratitude to Dr. Xiu-Guo Zhang and Dr. De-Wei Li for their critical review of the manuscript. The authors are grateful to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) (Project 88881.062172/2014-01) and Programa Ciência sem Fronteiras for financial support. RFCR is grateful to the Cuban Ministry of Agriculture and Programa de Salud Animal y Vegetal (Project P131LH003033) for facilities. We acknowledge the facilities provided by Dr. P.M. Kirk and Drs. V. Robert and K. Bensch, through the Index Fungorum and MycoBank websites. Dr. Lorelei Norvell's editorial review and Dr. Shaun Pennycook's nomenclature review are greatly appreciated.

Literature cited

- Carmichael JW. 1955. Lacto-fuchsin: a new medium for mounting fungi. *Mycologia* 47: 611.
- Castañeda-Ruiz RF. 1987. Fungi cubenses II. La Habana, Cuba. 22 p.
- Castañeda-Ruiz RF, Heredia G, Reyes M, Arias RM, Decock C. 2001. A revision of the genus *Pseudospiropes* and some new taxa. *Cryptogamie, Mycologie* 22: 3–18. [https://doi.org/10.1016/S0181-1584\(01\)01057-0](https://doi.org/10.1016/S0181-1584(01)01057-0)
- Castañeda-Ruiz RF, Guarro J, Velásquez-Noa S, Gené J. 2003. A new species of *Minimelanolocus* and some hyphomycete records from rain forests in Brazil. *Mycotaxon* 85: 231–239.
- Castañeda-Ruiz RF, Heredia G, Gusmão LFP, Li DW. 2016. Fungal diversity of Central and South America. 197–217, in: DW Li (ed.). *Biology of Microfungi*. Springer International Publishing. https://doi.org/10.1007/978-3-319-29137-6_9
- Fiuza PO, Conceição LB, Marques MFO, Gusmão LFP, Rafael F, Castañeda-Ruiz RF. 2017. *Dictyotrichocladium aquaticum* gen. & sp. nov. and *Minimelanolocus aquatilis* sp. nov. from freshwater in Brazil's semiarid region. *Mycotaxon* 132(2): 433–440. <https://doi.org/10.5248/132.433>.
- Heredia G, Arias RM, Castañeda-Ruiz RF, Minter DW. 2014. New species of *Lobatopedis* and *Minimelanolocus* (anamorphic fungi) from a Mexican cloud forest. *Nova Hedwigia* 98: 31–40. <https://doi.org/10.1127/0029-5035/2013/0146>
- Hernández-Restrepo M, Castañeda-Ruiz RF, Gené J, Guarro J, Minter DW, Stadler M. 2012. Microfungi from Portugal: *Minimelanolocus manifestus* sp. nov. and *Vermiculariopsiella pediculata* comb. nov. *Mycotaxon* 122: 135–143. <https://doi.org/10.5248/122.135>

- Liu XY, Udayanga D, Luo ZL, Chen LJ, Zhou DQ, Hyde KD. 2015. Backbone tree for *Chaetothyriales* with four new species of *Minimelanolocus* from aquatic habitats. *Fungal Biology* 119: 1046–1062. <https://doi.org/10.1016/j.funbio.2015.08.005>
- Ma J, Zhang YD, Ma LG, Zhang XG. 2008. Two new species of the *Minimelanolocus* in China. *Mycotaxon* 104: 147–151.
- Ma J, Zhang YD, Ma LG, Zhang XG. 2011. Two new *Minimelanolocus* species from southern China. *Mycotaxon* 117: 131–135. <https://doi.org/10.5248/117.131>
- Matsushima T. 1983. Matsushima mycological memoirs, no. 3. 90 p.
- Santa Izabel TS, Santos DS, Almeida DAC de, Gusmão LFP. 2011. Fungos conidiais do bioma Caatinga II. Novos registros para o continente americano, Neotrópico, América do Sul e Brasil. *Rodriguésia* 62(2): 229–240. <https://doi.org/10.1590/2175-7860201162201>
- Tian Q, Doilom M, Luo ZL, Chomnunti P, Bhat JD, Xu JC, Hyde KD. 2016. Introducing *Melanoctona tectonae* gen. et sp. nov. and *Minimelanolocus yunnanensis* sp. nov. (*Herpotrichiellaceae*, *Chaetothyriales*). *Cryptogamie, Mycologie* 37: 477–492. <https://doi.org/10.7872/crym/v37.iss4.2016.477>
- Xia JW, Ma LG, Castañeda-Ruiz RF, Zhang XG. 2014. *Minimelanolocus bicolorata* sp. nov., *Paradendryphiopsis elegans* sp. nov. and *Corynesporella bannaense* sp. nov. from southern China. *Mycoscience* 55: 299–307. <https://doi.org/10.1016/j.myc.2013.11.003>
- Zhang K, Fu HB, Zhang XG. 2009. Taxonomic studies of *Minimelanolocus* from Yunnan, China. *Mycotaxon* 109: 95–101. <https://doi.org/10.5248/109.95>
- Zhang YD, Ma J, Ma LG, Zhang XG. 2010. A new species of *Minimelanolocus* from Fujian, China. *Mycotaxon* 114: 373–376. <https://doi.org/10.5248/114.373>